

1-1-2000

Productivity Control of Fine Particle Transport to Equatorial Pacific Sediment

Ellen Thomas

Wesleyan University, ethomas@wesleyan.edu

Follow this and additional works at: <http://wescholar.wesleyan.edu/div3facpubs>

Recommended Citation

Thomas, Ellen, "Productivity Control of Fine Particle Transport to Equatorial Pacific Sediment" (2000). *Division III Faculty Publications*. Paper 109.

<http://wescholar.wesleyan.edu/div3facpubs/109>

This Article is brought to you for free and open access by the Natural Sciences and Mathematics at WesScholar. It has been accepted for inclusion in Division III Faculty Publications by an authorized administrator of WesScholar. For more information, please contact dschnaidt@wesleyan.edu, ljohnson@wesleyan.edu.

Productivity control of fine particle transport to equatorial Pacific sediment

E. Thomas¹ and K. K. Turekian

Center for the Study of Global Change, Department of Geology and Geophysics, Yale University, New Haven, Connecticut

K.-Y. Wei

Department of Geology, National Taiwan University, Taipei

Abstract. Accumulation rates of ³He (from cosmic dust), ²³⁰Th (produced in the water column), barite (produced in the water column during decay of organic matter), and Fe and Ti (arriving with wind-borne dust) all are positively correlated in an equatorial Pacific core (TT013-PC72; 01.1°N, 139.4°W; water depth 4298 m). These accumulation rates are also positively correlated with the accumulation rates of noncarbonate material. They are not significantly correlated to the mass accumulation rate of carbonate, which makes up the bulk of the sediment. The fluctuations in accumulation rates of these various components from different sources thus must result from variations in some process within the oceans and not from variations in their original sources. Sediment focusing by oceanic bottom currents has been proposed as this process [Marcantonio *et al.*, 1996]. We argue that the variations in the accumulation rates of all these components are dominantly linked to changes in productivity and particle scavenging (³He, ²³⁰Th, Fe, Ti) by fresh phytoplankton detritus (which delivers Ba upon its decay) in the equatorial Pacific upwelling region. We speculate that as equatorial Pacific productivity is a major component of global oceanic productivity, its variations over time might be reflected in variations in atmospheric levels of methanesulfonic acid (an atmospheric reaction product of dimethyl sulfide, which is produced by oceanic phytoplankton) and recorded in Antarctic ice cores.

1. Introduction

The eastern equatorial Pacific Ocean has been speculated to have a major influence on global climate [e.g., Cane, 1998]. New productivity in that region is an important component of global oceanic productivity [e.g., Archer and Maier-Reimer, 1994; Shimmiel and Jahnke, 1994]. Nutrient- and CO₂-rich waters well up in this region and are the largest natural source of CO₂ to the atmosphere because only a small fraction of the upwelled CO₂ is recycled biologically [Murray *et al.*, 1994]. Fluctuations in productivity and burial efficiency of the organic matter in this region on short (annual) and longer (glacial-interglacial) timescales thus have the potential to have a major impact on the global carbon cycle [e.g., Hansell *et al.*, 1997].

The sedimentary record of carbonate sedimentation over the last several glacial cycles in the equatorial Pacific has been studied for more than 45 years [e.g., Arrhenius, 1952; Hays *et al.*, 1969; Thompson and Saito, 1974; Shackleton and Opdyke, 1976; Valencia, 1977; Chuey *et al.*, 1987; Farrell and Prell, 1989; Rea *et al.*, 1991; Lyle *et al.*, 1988, 1992], but there is no agreement on its interpretation [e.g., Wei *et al.*, 1995; LaMontagne *et al.*, 1996]. Specifically, there is no agreement whether the fluctuations in carbonate percentage and carbonate accumulation rates reflect fluctuations in primary productivity [Archer,

1991], in dissolution [Farrell and Prell, 1989], or in sediment focusing by bottom currents [Marcantonio *et al.*, 1996]. If these fluctuations are driven by dissolution, there is no agreement on the question whether the fluctuations in dissolution are caused mainly by changes in deep-water corrosiveness, thus in deep-water circulation patterns [Farrell and Prell, 1989], or by changes in supply of organic carbon (thus productivity) [e.g., Archer, 1991; Hagemberg *et al.*, 1995].

If productivity fluctuations, such as the productivity increase by a factor of 2 postulated to have occurred during the last glacial at low latitudes [e.g., Pedersen, 1983; Herguera and Berger, 1991; Murray *et al.*, 1993; Paytan *et al.*, 1996], caused the carbonate fluctuations, we do not know what caused the changes in productivity. Different authors have invoked increased rates of upwelling [e.g., Berger and Wefer, 1991], increased rates of supply of limiting micronutrients such as iron [Coale *et al.*, 1996], or more complex processes involving nitrate utilization [Farrell *et al.*, 1995; Falkowski, 1997]. It is not clear whether any proxy (e.g., concentrations or mass accumulation rates of barite, opal, organic carbon, CaCO₃, or Al/Ti) reliably mirrors long-term productivity or at least delivery of organically produced material to the seafloor [e.g., McCorkle *et al.*, 1994; Pisias *et al.*, 1995], although primary production and particulate organic flux presently are well correlated in the equatorial Pacific [e.g., Betzer *et al.*, 1984; Smith *et al.*, 1997].

We evaluate published data on an equatorial Pacific core and suggest a mechanism (marine productivity and particle scavenging) by which all these different data sets can be explained in an internally consistent manner. We add a

¹Also at Department of Earth and Environmental Sciences, Wesleyan University, Middletown, Connecticut.

Table 1. Age Model of Core TT013-PC72^a

Isotope Event	Boundary of Stages	Age, kyr ^b	Depth in PC72, cm
2.0	1/2	12	35–40
3.0	2/3	24	75–80
4.0	3/4	59	135–140
5.0	4/5	71	165–170
6.0	5/6	128	235–240
7.0	6/7	186	368–373
8.0	7/8	245	448–453
9.0	8/9	303	543–548
10.0	9/10	339	604–609
11.0	10/11	362	638–644
12.0	11/12	423	742–747

^a From Murray *et al.* [1995] and Marcantonio *et al.* [1996].

^b Age (kyr) after Imbrie *et al.* [1984], names of isotope events and stage boundaries as listed by Prell *et al.* [1986].

speculation over possible teleconnections between equatorial Pacific productivity and deposition of methanesulfonic acid (MSA) in the Antarctic ice cap.




Figure 1a. Core TT013-PC72. Age model derived from correlation of benthic oxygen isotope data [Marcantonio *et al.*, 1996; A. Mix, personal communication, 1997] to the SPECMAP stack [Imbrie *et al.*, 1984]; gray bars indicate glacial isotope stages 2 through 12. Accumulation rate of carbonate [Murray *et al.*, 1995].

2. Data Compared

Core TT013-PC72 (140°W, 0.1°N; water depth 4298 m) was recovered as part of the equatorial Pacific Joint Global Ocean Flux Studies (JGOFS). An age model for its sediments was derived from $\delta^{18}\text{O}$ analysis of the benthic foraminifer *Cibicides wuellerstorfi* and correlation to the SPECMAP stack [Murray *et al.*, 1995; Marcantonio *et al.*, 1996; A. Mix, personal communication, 1997]. The benthic foraminiferal oxygen isotope record for core TT013-PC72 closely resembles the SPECMAP stacked record, with many of the subevents well represented. The depth of the stage boundaries is close to these in the late Quaternary reference section [Prell *et al.*, 1986] (Table 1). Numerical ages of the samples and accumulation rates can thus be estimated with confidence within the SPECMAP age model.

There is a strong positive correlation between accumulation rates of Fe and Ti (wind-borne dust), ^{230}Th (produced in the water column by radio-isotopic decay), ^3He (carried by interplanetary dust particles), and barite (produced by decay in association with organic matter), as can be seen by comparing data by Murray *et al.* [1995], Marcantonio *et al.* [1995, 1996], and Paytan *et al.* [1996] (Figure 1a–j). All these accumulation rates are significantly and positively correlated to each other. There is no simple, significant correlation between these accumulation rates and the oxygen isotopic record of benthic foraminifera, i.e., between these rates and global ice volume (Figure 2). The accumulation rates of these sediment compo-




Figure 1b. Same as Figure 1a, but for accumulation rate of noncarbonate [Murray *et al.*, 1995].




Figure 1c. Same as Figure 1a, but for oxygen isotope data (*Cibicidoides wuellerstorfi* [Marcantonio *et al.*, 1996; A. Mix, personal communication, 1997]).

nents [Murray *et al.*, 1995] are not significantly correlated to the accumulation rate of calcium carbonate, which makes up 60–90% of the sediment (Figure 3).

3. Discussion

The positive correlation between accumulation rates of various components that have different sources indicates that all these particles (wind-borne, cosmic dust, biogenic) were concentrated within the water column and that the fluctuations in their accumulation rates must have been caused by a common concentrating process within the oceans. Fluctuations in accumulation rates of Fe, for example, may not be caused directly by fluctuations in wind-borne dust concentrations (in contrast to Murray *et al.* [1995]). Fluctuations in the accumulation rate of ^3He are likewise probably not caused by fluctuations in the influx of cosmic dust (in contrast to Farley and Patterson [1995]).

We argue that sediment focusing may very well be a factor but that it cannot explain the correlations by itself alone: if there had been simple sediment focusing, there should have been a correlation between the accumulation rate of the non-carbonate components and that of bulk sediment. There is no such correlation (Figure 3) because there is no correlation

between the accumulation rates of these noncarbonate components and that of CaCO_3 , which determines the bulk accumulation rate. It is not probable that size-dependent winnowing only could have caused the differences in CaCO_3 and non- CaCO_3 accumulation rates because a large part of the carbonate is derived from nanoplankton in the same size range as the fine noncarbonate particles.

Marcantonio *et al.* [1995] suggested that CaCO_3 dissolution caused this lack of correlation, because the peaks of ^3He and ^{230}Th accumulation occurred at minima in $\text{CaCO}_3\%$ (Figures 1a–j and 4). They thus implied that low $\text{CaCO}_3\%$ values are caused by dissolution. In order to evaluate whether low carbonate content is caused by dissolution, we need to compare the $\text{CaCO}_3\%$ data with an index of dissolution, such as the index for dissolution of planktonic foraminifera [e.g., Berger, 1973; Le and Shackleton, 1992; LaMontagne *et al.*, 1996]. If we do this for core TT013-PC72, we see that there is no significant correlation between the $\text{CaCO}_3\%$ and the mass accumulation rates of CaCO_3 and between $\text{CaCO}_3\%$ and indices of dissolution such as the fragmentation of planktonic foraminifera (Figure 5).

If we assume an initial sediment composition of 85% carbonate, 15% noncarbonate (as usually done for equatorial Pacific sediments from depths of around 4200 m [Pisias and Prell, 1985]), a change to a carbonate percentage of 60% (about the lowest value observed) would require dissolution of $\sim 75\%$ of the original carbonate [Dean *et al.*, 1981]. Such




Figure 1d. Same as Figure 1a, but for accumulation rate of Fe [Murray *et al.*, 1995].




Figure 1e. Same as Figure 1a, but for accumulation rate of Ti [Murray *et al.*, 1995].

severe dissolution does not agree with the observed indices of dissolution (Figure 5). Decoupling of $\text{CaCO}_3\%$ and dissolution intensity has been observed in many cores in the equatorial Pacific [e.g., Luz and Shackleton, 1975; Wei *et al.*, 1995; LaMontagne *et al.*, 1996].

We argue that the correlation between CaCO_3 minima and the excess deposition of ^3He and ^{230}Th can be seen as an indirect argument against sediment focusing as a cause of this excess deposition. The $\text{CaCO}_3\%$ minima have been widely correlated among many cores in the equatorial Pacific region [e.g., Arrhenius, 1952; Hays *et al.*, 1969; Luz and Shackleton, 1975; Farrell and Prell, 1989]. If these minima are linked to sediment focusing, the excess deposition resulting from the focusing must have also occurred throughout the eastern equatorial Pacific “sediment bulge,” which leaves the place of origin from where the focused sediment must have been removed far outside this bulge. Such an outside source appears more probable if we envisage the material to have been moved (at least in part) toward the “equatorial bulge” by lateral advection toward the equator in surface to upper intermediate waters rather than by bottom currents.

Marcantonio *et al.* [1996] favored sediment focusing by bottom currents as the mechanism and cited sediment trap data [Lao *et al.*, 1993; François *et al.*, 1990] to argue that organically produced particle scavenging could not be the cause of the correlation between all the accumulation rates. We agree

with François *et al.* [1990] that sediment focusing is the most important factor in equatorial Atlantic sediments from a non-upwelling region but consider these observations not relevant to the interpretation of carbonate sedimentation and dissolution in an upwelling region in the Pacific. In the equatorial Atlantic cores from nonupwelling regions, for instance, carbonate productivity was lower during the last glacial [François *et al.*, 1990], in contrast with most interpretations of the eastern equatorial upwelling region during the last glacial [e.g., Pedersen, 1983; Herguera and Berger, 1991; Murray *et al.*, 1993; Paytan *et al.*, 1996]. Others have argued that sediment trap data may not accurately reflect the spatial and temporal heterogeneity of the biogenic flux [Buesseler *et al.*, 1994; Boyd and Newton, 1997; Beaulieu and Smith, 1998; Shaw *et al.*, 1998].

JGOFS data along 140°W [Honjo *et al.*, 1995] demonstrate strong variability in space and time and demonstrate the occurrence of very short-term (less than 3 weeks) episodes of high-particle flux during the passage of tropical instability waves. We suggest specifically that the well-recognized overestimates of particle flux during low-productivity periods and underestimates at high-productivity periods may be more severe than suggested [François *et al.*, 1993; Buesseler, 1998]. Shaw *et al.* [1998], for instance, document that the accumulation rate of Th nuclides is underestimated by a factor of 10 using sediment trap data in the northeastern Pacific.




Figure 1f. Same as Figure 1a, but for the ^3He burial rate [Marcantonio *et al.*, 1996].




Figure 1g. Same as Figure 1a, but for accumulation rate of barite [Paytan *et al.*, 1996].

We also conclude that sediment accumulated during these high-flux events may make up a significant part of the total sediment deposited [Kemp *et al.*, 1995; Verity *et al.*, 1996; Shaw *et al.*, 1998]. On the basis of the above considerations we argue that fluctuations in the abundance of scavenging phytodetritus which is deposited rapidly to the seafloor may have caused the fluctuations in accumulation rates of all the various materials associated with fine, noncarbonate particles derived from different sources. JGOFS studies along 140°W document that presently, phytodetritus accumulates on the seafloor in a pattern reflecting overall productivity [Smith *et al.*, 1996; Barber *et al.*, 1996], with the highest accumulation rates close to the location of core TT013-PC72. This phytodetritus may largely be concentrated in dense mats containing needle-shaped diatoms [Kemp *et al.*, 1995], which are concentrated during the passage of tropical instability waves [e.g., Kemp, 1994; Yoder *et al.*, 1994; Smith *et al.*, 1996]. Such long waves occur seasonally, are weak during El Niño events [Feely *et al.*, 1994], but are strong just after its ending [Yoder *et al.*, 1994; Smith *et al.*, 1996].

In the present ocean, removal of Th isotopes from seawater column is strongly linked to biological activity [e.g., Lao *et al.*, 1993; Shaw *et al.*, 1998]. This phytodetrital scavenging is at present probably not a very important contributor to the sediment flux, but it might have been more important during earlier geological periods [e.g., Kemp, 1995]. We suggest that at times of higher equatorial Pacific upwelling rates, produc-

tivity was higher. At these times, phytoplankton blooms were more frequent and rapidly deposited phytodetrital matter contributed a larger fraction of the total sediment compared to the present [e.g., Ittekkot, 1993; Verity *et al.*, 1996; Buesseler, 1998; Shaw *et al.*, 1998]. We thus envisage that the strong enrichment in ^{230}Th (more than can be derived from the water column directly over the site of the core) as caused by a combination of high and fluctuating productivity, scavenging of elements by organically produced particles, and concentration of such particles from a fairly large region by passage of tropical instability waves.

Accumulation rates of the various fine-grain particles associated components (such as cosmic He) and scavenged elements (such as Ti and Al) can therefore be used as a proxy for delivery of organically produced material to the seafloor. In the equatorial Pacific the particulate organic flux is linked to primary productivity [Betzer *et al.*, 1984], and the vertical flux of biogenic particles has been shown to exert tight control on the nature and rates of benthic biological and chemical processes [Smith *et al.*, 1997]. We therefore can use the accumulation rates of particles that are indirectly (by scavenging of dissolved elements or small particles) associated with organic matter flux as a proxy for primary productivity.

This conclusion is supported by the positive correlation with the barite accumulation rate because barite forms in microenvironments within decaying organic matter [e.g., Dehairs *et al.*,




Figure 1h. Same as Figure 1a, but for burial rate of excess ^{230}Th [Marcantonio *et al.*, 1995, 1996].




Figure 1i. Same as Figure 1a, but for dissolution of carbonate in core TT013-PC72, using the percentage of whole planktonic foraminifera as an indicator (K. Y. Wei, unpublished data, 1999).

1980, 1990; Bishop, 1988]. Barite accumulation rates are thus directly related to the organic matter export flux [Paytan *et al.*, 1996] and to the accumulation rates of organic carbon in the sediments [Lyle *et al.*, 1988, 1992]. Further support comes from a comparison of the accumulation rates of various fine-grain particle associated components with the record of combustion oxygen demand (COD) over the last 400 kyr at eastern equatorial Pacific Ocean Drilling Program (ODP) site 849 at $0^{\circ}11'N$, $110^{\circ}34'W$ and a water depth of 3839 m [Perks, 1999]. This parameter provides a good approximation of the organic carbon content of sediments, which have such low-organic carbon content that it cannot be reliably measured. COD data [Perks, 1999] are available at higher time resolution than the data for core TT013-PC72, and we thus cannot compare the records sample by sample. There are, however, peaks in COD at site 849 and peaks in the fine-grain particle associated components in core TT013-PC72 at 20 kyr, 70 kyr, 150–185 kyr, 275 kyr, and 320 kyr.

This view implies that the calcium carbonate accumulation rate at the location of TT013-PC72 cannot be used as productivity proxy: at very high productivity the main primary producers are the Si-walled diatoms rather than the carbonate-walled nannoplankton [e.g., Dymond and Collier, 1988]. Calcite-secreting and silica-secreting plankton communities may alternate in dominating overall productivity [Lyle *et al.*, 1988]. In the equatorial Pacific, high productivity could be reflected by low values of $CaCO_3\%$, as a result of a combina-

tion of dilution by biogenic silica and increased dissolution caused by the larger supply of organic material [e.g., Theyer *et al.*, 1985; Hagelberg *et al.*, 1995]. The deposition of the various components linked to organic material does not necessarily imply high net accumulation rates of organic matter in the sediments because labile organic matter is quickly degraded on the seafloor by bacteria and protists, given an adequate oxygen supply [Gooday and Turley, 1990; Turley and Lochte, 1990; Poremba, 1994].

We do not know whether the overall biological composition of the bulk phytodetritus (thus its biogenic silica and organic matter content) would have been constant over time. Probably, it fluctuated on glacial-interglacial timescales with fluctuations in the biological components [e.g., Lyle *et al.*, 1988]. We can therefore not assume that rates of element concentrations within this phytodetritus have been constant over time. The ratio of scavenged material to organically produced material likewise may have fluctuated.

The peaks in accumulation rates of the above-mentioned indicators are to some extent but not completely related to glacial isotope stages: comparison of the records with the oxygen isotope data from core PC72 (Figure 1a) shows that they occurred in glacials 2, 4, 6, and 8 (but were much more short-lived than glacials 6 and 8) but also in interglacials 9 and 11, which are said to represent the warmest two interglacials [e.g., Hodell, 1993]. There is no peak in accumulation rate in biogenic Ba apparent in interglacial 9, but this is probably caused by a lack of data points over the core interval with the




Figure 1j. Same as Figure 1a, but for percentage of $CaCO_3$ in core TT013-PC72 [Murray *et al.*, 1995].




Figure 2. Comparison of accumulation rates of various sediment components and the benthic foraminiferal oxygen isotope data for the same samples. Accumulation rates after Murray *et al.*, [1995] and oxygen isotope values as given by Marcantonio *et al.* [1996] and A. Mix (personal communication, 1997). Here r^2 indicates value of correlation coefficient. (a) CaCO_3 , (b) non- CaCO_3 , (c) Fe, and (d) Ti.

peak in mass accumulation rates of the other productivity indicators.

The linkage between fluctuations in $\text{CaCO}_3\%$, CaCO_3 accumulation rates, and glacial-interglacial timing is complex [e.g., Luz and Shackleton, 1975; Farrell and Prell, 1989]. Productivity in tropical Pacific regions probably was not coupled to ice-volume fluctuations, the variance of which was dominated by the 100 kyr Milankovich cyclicity during the last 900 kyr (Figure 2). Tropical productivity fluctuated dominantly at the periodicity of precessional forcing (19–23 kyr) in the equatorial Indian Ocean [Beaufort *et al.*, 1997, 1999], western equatorial Pacific [Perks and Keeling, 1998; Cane, 1998], and eastern equatorial Pacific [Perks, 1999]. Perks' [1999] records of COD (a productivity proxy, see above) show peaks at times of peaks

in the mass accumulation rates of the productivity indicators observed in core TT013-PC72, although the time resolution of these records is not quite sufficient to resolve precessional periodicity.

Variability on timescales differing from the main glacial-interglacial 100 kyr periodicity in ice volume may explain that not all glacials at the location of TT013-PC72 were similar: during glacials 2, 4, and 6, peaks in carbonate and noncarbonate accumulation rates appear to be at least generally coeval, but during the earlier glacials and in interglacial 11, they are strongly decoupled. Carbonate mass accumulation rates are generally high in glacials but are not well correlated to the benthic foraminiferal oxygen isotope record on the same core (Figure 1a–j).




Figure 3. (a) Comparison of the accumulation rates of Fe and bulk sediment in core PC72 [Murray *et al.*, 1995]. (b) Comparison of bulk accumulation rates with accumulation rates of CaCO₃ and non-CaCO₃ [Murray *et al.*, 1995].

The correlation between CaCO₃% values and glacial-interglacial stages (Figure 1a–j) is even less precise. There is no clear correlation between CaCO₃% values and the transition between glacial and interglacial stages, in contrast with Broecker and Sanyal [1997]. These authors argued that dissolution events in core TT013-PC72 occurred at glacial/interglacial stage boundaries, but of the three events that they mentioned, only one (stage 7–6) occurred within a few thousand years of such a transition (Figure 1a–j). The earlier minima occur at about the middle of glacial stage 8 and interglacial stage 11; the latter of these can be correlated to the widely recognized carbonate minimum B9 [Hays *et al.*, 1969; Farrell and Prell, 1989].

Understanding the coupling and decoupling of the accumulation rates of carbonate and noncarbonate is of importance for the understanding of glacial/interglacial changes in equatorial Pacific

climate. It is evident that the last glacial was not typical of all earlier glacials (Figure 1a–j), possibly because the variability in precession and eccentricity bands interacts differently during different isotope stages [e.g., Beaufort *et al.*, 1999].

Equatorial Pacific productivity is a large component of global oceanic productivity [Barber and Chavez, 1987], and productivity fluctuations in eastern and western Pacific appear to be correlated [Perks, 1999]. Therefore we argue that equatorial Pacific productivity fluctuations could be seen as an important factor of global oceanic productivity. If this is correct, we might see an impact of variations in equatorial Pacific productivity in global records of methanesulfonic acid (MSA), a reaction product of dimethyl sulfide which is produced by phytoplankton. Dimethyl sulfide (DMS) and its reaction products are indeed seen to be enriched across the high-productivity




Figure 4. Comparison of the percentage of CaCO₃ [Murray *et al.*, 1995] and the accumulation rates of (a) Fe and (b) Ti.

tivity zone in the equatorial Pacific along 140°W [Andreae and Raemdonck, 1983; Charlson *et al.*, 1987], although Bates and Quinn [1997] observed no variations in the high DMS concentrations in surface waters of the equatorial Pacific during 1982–1996 El Niño cycles.

A relation between equatorial Pacific productivity and the Antarctic MSA record appears to exist in records over the last 150 kyr [LeGrand *et al.*, 1991], as seen from the general agreement in accumulation rate of biogenic Ba in TT013-PC72 with that of MSA in the Antarctic Vostok ice core (Figure 6). Specifically, the peak in equatorial productivity at ~ 70 kyr (glacial isotope stage 4) corresponds to a peak in Antarctic MSA.

We do not think that these high concentrations in MSA in Antarctic ice cores during glacial times were derived from productivity close to the Antarctic continent, although LeGrand and Feniet-Saigne [1991] linked variations in MSA accumulation rates in Antarctic snow layers to strong El Niño events. They argued that the enrichments in MSA were produced close to the Antarctic continent, possibly as a result of El Niño–Southern Oscillation linked variations in sea ice cover because in the present ocean, high levels of DMS are produced during the Southern Hemispheric summer by *Phaeocystis* [DiTullio *et al.*, 1998; Kettle *et al.*, 1999]. Even during cold periods in the Holocene, however, productivity was low in the Antarctic [e.g., Leventer *et al.*, 1996], and Antarctic productivity during the Last Glacial Maximum was depressed [e.g., François *et al.*, 1993; Kumar *et al.*, 1993]. We therefore argue that the shelf regions where relatively high productivity occurs in the Southern Hemisphere summer would have much more extensive ice cover during glacials, limiting seasonal phytodetrital productivity. The elevated MSA levels thus would have to be transported from a place with elevated productivity during glacial periods, such as the equatorial Pacific, although we do not know whether the apparent correlation is due primarily to




Figure 5. Comparison of the percentage CaCO_3 [Murray *et al.*, 1995] with the percentage of whole foraminifera, an indicator of severity of dissolution.




Figure 6. Accumulation rates of Ba in equatorial Pacific core TT013-PC72 (open symbols [Murray *et al.*, 1995]) compared to accumulation rates of MSA in the Antarctic Vostok ice core (solid symbols [LeGrand *et al.*, 1991]). The correlation was made using the depths provided by LeGrand *et al.* [1991] for the Vostok ice core and calculating numerical ages according to Sowers *et al.* [1993].

changes in equatorial Pacific productivity patterns or to changes in the efficiency of atmospheric transport to Antarctica.

4. Conclusions

Accumulation rates of sediment components derived from interplanetary dust (^3He), from radioactive decay in the water column (^{230}Th), from biological activity in the water column (barite), and from wind-blown dust (Fe, Ti) are strongly correlated in an equatorial Pacific carbonate core. We suggest that this strong correlation results dominantly from concentration of all these components by particle scavenging of organically produced particulate matter (^3He , ^{230}Th , Fe, Ti) and decomposition of the organic matter (Ba). Both productivity and efficiency of scavenging may play a role. The accumulation rates of these components thus are proxies for productivity.

We speculate that global oceanic productivity is strongly influenced by equatorial Pacific productivity and that this global signal might be present in the MSA records in Antarctic ice cores.

Acknowledgments. E. Thomas acknowledges partial funding of research by the 1996 Storrs-Cole Award (GSA) and thanks John King for providing samples from core TT013-PC72. The research was also supported by the Global Change Program of the National Science Foundation administered by the Earth Sciences Division. We thank Bob Anderson, Michael Bender, and Rick Murray for reviews.

References

- Andreae, M. O., and H. Raemdonck, Dimethylsulfide in the surface ocean and the marine atmosphere: A global view, *Science*, **221**, 744–747, 1983.
- Archer, D. E., Equatorial Pacific calcite preservation cycles: Production or dissolution?, *Paleoceanography*, **6**, 561–571, 1991.
- Archer, D., and E. Maier-Reimer, Effect of deep-sea sedimentary calcite preservation on atmospheric CO₂ concentration, *Nature*, **367**, 260–263, 1994.
- Arrhenius, G., Sediment cores from the east Pacific: Properties of the sediments, *Swedish Deep Sea Exped., Rep. 1951–1952*, 1–227, 1952.
- Barber, R. T., and F. P. Chavez, Regulation of primary productivity in the equatorial Pacific Ocean, *Limnol. Oceanogr.*, **36**, 1803–1815, 1987.
- Barber, R. T., M. P. Sanderson, S. T. Lindley, F. Chai, C. C. Trees, D. G. Foley, and F. P. Chavez, Primary productivity and its regulation in the equatorial Pacific during and following the 1991–1992 El Niño, *Deep Sea Res., Part II*, **43**, 933–969, 1996.
- Bates, T. S., and P. K. Quinn, Dimethylsulfide (DMS) in the equatorial Pacific Ocean (1982 to 1996): Evidence of a climate feedback?, *Geophys. Res. Lett.*, **24**, 861–864, 1997.
- Beaufort, L., Y. Lancelot, P. Camberlin, O. Cayre, E. Vincent, F. Bassinot, and L. Labeyrie, Insolation cycles as a major control of the equatorial Indian Ocean primary productivity, *Science*, **278**, 1451–1454, 1997.
- Beaufort, L., F. Bassinot, and E. Vincent, Primary production response to orbitally induced variations of the Southern Oscillation in the equatorial Indian Ocean, in *Reconstructing Ocean History: A Window Into the Future*, edited by F. Abrantes and A. Mix, pp. 245–271, Plenum, New York, 1999.
- Beaulieu, S. E., and K. L. Smith Jr., Phytodetritus entering the benthic boundary layer and aggregated on the sea floor in the abyssal NE Pacific: Macro- and microscopic comparison, *Deep Sea Res., Part II*, **45**, 781–815, 1998.
- Berger, W. H., Deep-sea carbonates: Pleistocene dissolution cycles, *J. Foraminiferal Res.*, **3**, 187–195, 1973.
- Berger, W. H., and G. Wefer, Productivity of the glacial ocean: Discussion of the iron hypothesis, *Limnol. Oceanogr.*, **36**, 1899–1918, 1991.
- Betzler, P. R., W. J. Showers, E. A. Laws, C. D. Winn, G. R. DiTullio, and P. M. Kroopnick, Primary productivity and particle fluxes on a transect of the equator at 135°W in the Pacific Ocean, *Deep Sea Res., Part A*, **31**, 1–11, 1984.
- Bishop, J. K. B., The barite-opal-organic carbon association in oceanic particulate matter, *Nature*, **332**, 341–343, 1988.
- Boyd, P., and P. Newton, Measuring biogenic carbon flux in the ocean, *Science*, **275**, 554, 1997.
- Broecker, W. S., and A. Sanyal, Magnitude of the CaCO₃ dissolution events marking the onset of times of glaciation, *Paleoceanography*, **12**, 530–532, 1997.
- Buesseler, K. O., The decoupling of production and particulate export in the surface ocean, *Global Biogeochem. Cycles*, **12**, 297–310, 1998.
- Buesseler, K. O., A. F. Michaels, D. A. Siegel, and A. H. Knap, A three dimensional time-dependent approach to calibrating sediment fluxes, *Global Biogeochem. Cycles*, **8**, 179–193, 1994.
- Cane, M. A., A role for the Tropical Pacific, *Science*, **282**, 59–61, 1998.
- Charlson, R. J., J. E. Lovelock, M. O. Andreae, and S. G. Warren, Oceanic phytoplankton, atmospheric sulfur cloud albedo and climate, *Nature*, **326**, 655–661, 1987.
- Chuey, J. M., D. K. Rea, and N. G. Pisias, Late Pleistocene paleoclimatology of the central equatorial Pacific: A quantitative record of eolian and carbonate deposition, *Quat. Res.*, **28**, 323–339, 1987.
- Coale, K. H., et al., A massive phytoplankton bloom induced by an ecosystem-scale iron fertilization experiment in the Pacific Ocean, *Nature*, **383**, 495–501, 1996.
- Dean, W. E., J. V. Gardner, and P. Cepek, Tertiary carbonate dissolution cycles on the Sierra Leone Rise eastern equatorial Atlantic Ocean, *Mar. Geol.*, **39**, 81–101, 1981.
- Dehairs, F., R. Chesselet, and J. Jedwab, Discrete suspended particles of barite and the barium cycle in the oceans, *Earth Planet. Sci. Lett.*, **49**, 528–550, 1980.
- Dehairs, F., L. Goeyens, N. Stroobants, P. Bernard, C. Goyet, A. Poisson, and R. Chesselet, On suspended barite and the oxygen minimum in the Southern Ocean, *Global Biogeochem. Cycles*, **4**, 85–102, 1990.
- DiTullio, G. R., D. L. Garrison, and S. Matho, Dimethylsulfoniopropionate in sea ice algae from the Ross Sea polynya, *Antarct. Res. Ser.*, **73**, 139–146, 1998.
- Dymond, J., and R. Collier, Biogenic particle fluxes in the equatorial Pacific: Evidence for both high and low productivity during the 1982–1983 El Niño, *Global Biogeochem. Cycles*, **2**, 129–137, 1988.
- Falkowski, P. G., Evolution of the nitrogen cycle and its influence on the biological sequestration of CO₂ in the ocean, *Nature*, **387**, 272–275, 1997.
- Farley, K., and D. B. Patterson, A 100 kyr periodicity in the flux of extraterrestrial ³He to the sea floor, *Nature*, **378**, 600–603, 1995.
- Farrell, J. W., and W. L. Prell, Climate change and CaCO₃ preservation: An 800,000 year bathymetric reconstruction from the central equatorial Pacific Ocean, *Paleoceanography*, **4**, 447–466, 1989.
- Farrell, J. W., T. F. Pedersen, S. E. Calvert, and B. Nielsen, Glacial-interglacial changes in nutrient utilization in the equatorial Pacific Ocean, *Nature*, **377**, 514–517, 1995.
- Feely, R. A., R. Wanninkhof, C. E. Costa, M. J. McPhaden, R. H. Byrne, F. J. Millero, F. P. Chavez, T. Clayton, D. M. Campbell, and P. P. Murphy, The effect of tropical instability waves on CO₂ species distributions along the equator in the eastern equatorial Pacific during the 1992 El Niño event, *Geophys. Res. Lett.*, **21**, 277–280, 1994.
- François, R., M. P. Bacon, and D. O. Suman, Thorium-230 profiling in deep-sea sediments; High resolution records of flux and dissolution of carbonate in the equatorial Atlantic during the last 24,000 years, *Paleoceanography*, **5**, 761–787, 1990.
- François, R., M. P. Bacon, M. A. Altabet, and L. D. Labeyrie, Glacial/interglacial changes in sediment rain rate in the SW Indian sector of the subantarctic waters as recorded by ²³⁰Th, ²³¹Pa, U, and ¹⁵N, *Paleoceanography*, **8**, 611–629, 1993.
- Gooday, A. J., and C. M. Turley, Responses by benthic organisms to inputs of organic material to the ocean floor: A review, *Philos. Trans. R. Soc. London, Ser. A*, **331**, 119–138, 1990.
- Hagelberg, T., N. G. Pisias, L. A. Mayer, N. J. Shackleton, and A. C. Mix, Spatial and temporal variability of late Neogene equatorial Pacific carbonate: Leg 138, *Proc. Ocean Drill. Program Sci. Results*, **138**, 321–336, 1995.
- Hansell, D. A., N. R. Bates, and C. A. Carlson, Predominance of vertical loss of carbon from surface waters of the equatorial Pacific Ocean, *Nature*, **386**, 59–81, 1997.
- Hays, J. D., T. Saito, N. D. Opdyke, and L. H. Burckle, Pliocene-Pleistocene sediments of the equatorial Pacific: Their paleomagnetic, biostratigraphic and climatic record, *Geol. Soc. Am. Bull.*, **80**, 1481–1514, 1969.
- Herguera, J. C., and W. H. Berger, Paleoproductivity from benthic foraminifera abundance: Glacial to postglacial change in the west equatorial Pacific, *Geology*, **19**, 1173–1176, 1991.
- Hodell, D. A., Late Pleistocene paleoceanography of the South Atlantic section of the Southern Ocean: ODP Hole 704A, *Paleoceanography*, **8**, 47–67, 1993.
- Honjo, S., J. Dymond, R. Collier, and S. J. Manganini, Export production of particles to the interior of the equatorial Pacific Ocean during the 1992 EqPac experiment, *Deep Sea Res., Part II*, **44**, 831–870, 1995.
- Imbrie, J. J., J. D. Hays, D. G. Martinson, A. McIntyre, A. C. Mix, J. J. Morley, N. G. Pisias, W. L. Prell, and N. J. Shackleton, The orbital theory of Pleistocene climate: Support from a revised chronology of the ¹⁸O record, in *Milankovitch and Climate*, edited by A. L. Berger et al., pp. 269–305, D. Riedel, Norwell, Mass., 1984.
- Ittekkot, V., The abiotically driven biological pump in the ocean and short-term fluctuations in atmospheric CO₂ contents, *Global Planet. Change*, **8**, 17–25, 1993.
- Kemp, A. E. S., Laminated sediments from coastal and open ocean upwelling systems: What variability do they record?, in *Upwelling in the Ocean:*

- Modern Processes and Ancient Records*, edited by C. P. Summerhayes et al., pp. 239–257, John Wiley, New York, 1994.
- Kemp, A. E. S., J. G. Baldauf, and R. B. Pearce, Origins and paleoceanographic significance of laminated diatom ooze from the eastern equatorial Pacific, *Proc. Ocean Drill. Program*, 138, 641–663, 1995.
- Kettle, A. J., et al., A global database of sea surface dimethylsulfide and a procedure to predict sea surface DMS as a function of latitude, longitude and month, *Global Biogeochem. Cycles*, 13, 399–444, 1999.
- Kumar, N., R. Gwiazda, R. F. Anderson, and P. N. Froelich, $^{231}\text{Pa}/^{230}\text{Th}$ ratios in sediments as a proxy for past changes in Southern Ocean productivity, *Nature*, 372, 45–48, 1993.
- LaMontagne, R. W., R. W. Murray, K.-Y. Wei, M. Leinen, and C. H. Wang, Decoupling of carbonate preservation, carbonate concentration and biogenic accumulation: A 400 kyr record from the central equatorial Pacific Ocean, *Paleoceanography*, 11, 553–562, 1996.
- Lao, Y., R. F. Anderson, W. S. Broecker, H. Hofmann, and W. Wolfli, Particulate fluxes of ^{230}Th , ^{231}Pa and ^{10}Be in the northeastern Pacific Ocean, *Geochim. Cosmochim. Acta*, 57, 205–217, 1993.
- Le, J., and N. J. Shackleton, Carbonate dissolution fluctuations in the western equatorial Pacific during the late Quaternary, *Paleoceanography*, 7, 21–42, 1992.
- LeGrand, M., and C. Feniet-Saigne, Methanesulfonic acid in South Polar snow layers: A record of strong El Niño?, *Geophys. Res. Lett.*, 18, 187–190, 1991.
- LeGrand, M., C. Feniet-Saigne, E. S. Saltzman, C. Germain, I. Barkov, and V. N. Petrov, Ice-core record of oceanic emissions of dimethylsulfide during the last climate cycle, *Nature*, 350, 144–146, 1991.
- Leventer, A., E. W. Domack, S. E. Ishman, S. Brachfeld, C. E. McClennen, and P. Manley, Productivity cycles of 200–300 years in the Antarctic Peninsula region: Understanding linkages along the Sun, atmosphere, oceans, sea ice and biota, *Geol. Soc. Am. Bull.*, 108, 1626–1644, 1996.
- Luz, B., and N. J. Shackleton, CaCO_3 solution in the tropical east Pacific during the past 130,000 years, *Spec. Publ. Cushman Found. Foraminiferal Res.*, 13, 142–150, 1975.
- Lyle, M., D. W. Murray, B. P. Finney, J. Dymond, J. M. Robbins, and K. Brooksforce, The record of late Pleistocene biogenic sedimentation in the eastern tropical Pacific Ocean, *Paleoceanography*, 3, 39–59, 1988.
- Lyle, M., F. G. Prahl, and M. A. Sparrow, upwelling and productivity changes inferred from a temperature record in the central equatorial Pacific, *Nature*, 355, 812–815, 1992.
- Marcantonio, F., N. Kumar, M. Stute, R. F. Anderson, M. A. Seidl, P. Schlosser, and A. Mix, A comparative study of accumulation rates derived by He and Th isotope analysis of marine sediments, *Earth Planet. Sci. Lett.*, 133, 549–555, 1995.
- Marcantonio, F., R. F. Anderson, M. Stute, N. Kumar, P. Schlosser, and A. Mix, Extraterrestrial ^3He as a constant-flux tracer for paleoceanographic studies, *Nature*, 383, 705–707, 1996.
- McCorkle, D. C., D. H. Veeh, and D. T. Heggie, Glacial-Holocene paleo-productivity off western Australia: A comparison of proxy records, in *Carbon Cycling in the Glacial Ocean: Constraints on the Ocean's Role in Global Change*, edited by R. Zahn et al., *NATO ASI Ser., Ser. I*, 17, 443–479, 1994.
- Murray, J. W., R. T. Barber, M. R. Roman, M. P. Bacon, and R. A. Feely, Physical and biological controls on carbon cycling in the equatorial Pacific, *Science*, 266, 58–65, 1994.
- Murray, R. W., M. Leinen, and A. Isern, Biogenic flux of Al to sediment in the central equatorial Pacific Ocean: Evidence for increased productivity during glacial periods, *Paleoceanography*, 6, 651–669, 1993.
- Murray, R. W., M. Leinen, D. W. Murray, A. C. Mix, and C. W. Knowlton, Terrigenous Fe input and biogenic sedimentation in the glacial and interglacial equatorial Pacific Ocean, *Global Biogeochem. Cycles*, 9, 667–684, 1995.
- Paytan, A., M. Kastner, and F. P. Chavez, Glacial to interglacial fluctuations in productivity in the equatorial Pacific as indicated by marine barite, *Science*, 274, 1355–1357, 1996.
- Pedersen, T. F., Increased productivity in the eastern equatorial Pacific during the last glacial maximum, *Geology*, 11, 16–19, 1983.
- Perks, H., Late Pleistocene precessional forcing in the eastern and western equatorial Pacific, *JOI/USSAC News*, 12, 15, 1999.
- Perks, H. M., and R. F. Keeling, A 400 kyr record of combustion oxygen demand in the western equatorial Pacific: Evidence for precessionally forced climate response, *Paleoceanography*, 13, 63–69, 1998.
- Pisias, N. G., and W. L. Prell, Changes in calcium carbonate accumulation in the equatorial Pacific during the late Cenozoic: Evidence from HPC Site 572, in *The Carbon Cycle and Atmospheric CO_2 : Natural Variations Archaean to Present*, *Geophys. Monogr. Ser.*, vol. 32, edited by E. T. Sundquist and W. S. Broecker, pp. 443–454, AGU, Washington, D. C., 1985.
- Pisias, N. G., et al., Marine Aspects of Earth System History (MESH) Proxy Development Workshop, Proxy Workshop Rep., 75 pp., Corvallis, Oreg., Jan. 1995.
- Poremba, K., Simulated degradation of phytodetritus in deep-sea sediments of the NE Atlantic ($47^\circ\text{N}, 19^\circ\text{W}$), *Mar. Ecol. Prog. Ser.*, 105, 291–299, 1994.
- Prell, W., J. Imbrie, D. G. Martinson, J. J. Morley, N. G. Pisias, N. J. Shackleton, and H. F. Streeter, Graphic correlation of oxygen isotope stratigraphy application to the late Quaternary, *Paleoceanography*, 1, 137–162, 1986.
- Rea, D. K., N. G. Pisias, and T. Newberry, Late Pleistocene paleoclimatology of the central equatorial Pacific: Flux patterns of biogenic sediments, *Paleoceanography*, 6, 227–244, 1991.
- Shackleton, N. J., and N. D. Opdyke, Oxygen-isotope and paleomagnetic stratigraphy of Pacific core V28-239 Late Pliocene to Latest Pleistocene, *Mem. Geol. Soc. Am.*, 145, 449–464, 1976.
- Shaw, T. J., J. M. Smoak, and L. Lauerma, Scavenging of ex^{234}Th , ex^{230}Th , and ex^{210}Pb by particulate matter in the water column of the California Continental Margin, *Deep Sea Res., Part II*, 45, 763–779, 1998.
- Shimmield, G. B., and R. A. Jahnke, Particle flux and its conversion to the sediment record: Open ocean upwelling systems, in *Upwelling in the Ocean: Modern Processes and Ancient Records*, edited by C. P. Summerhayes, et al., pp. 171–191, John Wiley, New York, 1994.
- Smith, C. R., D. J. Hoover, S. E. Doan, R. H. Pope, D. J. Demaster, F. C. Dobbs, and M. Altabet, Phytodetritus at the abyssal seafloor across 10° of latitude in the central equatorial Pacific ocean, *Deep Sea Res. Part II*, 43, 1309–1338, 1996.
- Smith, C. R., W. Berelson, D. J. DeMaster, F. C. Dobbs, D. Hammond, D. J. Hoover, R. H. Pope, and M. Stephens, Latitudinal variations in benthic processes in the abyssal equatorial Pacific: Control by biogenic particle flux, *Deep Sea Res. Part II*, 44, 2295–2317, 1997.
- Sowers, T., M. Bender, L. Labeyrie, D. Martinson, J. Jouzel, D. Raynaud, J.-J. Pichon, and Y. S. Korotkevich, A 135,000-year Vostok-SPECMAP common temporal framework, *Paleoceanography*, 8, 737–766, 1993.
- Theyer, F., L. A. Mayer, J. A. Barron, and E. Thomas, The equatorial Pacific high-productivity belt: Elements for a synthesis of DSDP Leg 85 Results, *Initial Rep. Deep Sea Drill. Proj.*, 85, 971–985, 1985.
- Thompson, P. R., and T. Saito, Pacific Pleistocene sediments: Planktonic foraminifera dissolution cycles and geochronology, *Geology*, 2, 333–335, 1974.
- Turley, C. M., and K. Lochte, Microbial response to the input of fresh detritus to the deep-sea bed, *Palaeogeogr. Palaeoclim. Palaeoecol.*, 89, 3–23, 1990.
- Valencia, M., Pacific Pleistocene paleoclimatic stratigraphies: A comparative analysis of results, *Quat. Res.*, 8, 339–354, 1977.
- Verity, P. G., D. K. Stoecker, M. E. Sierack, and J. R. Nelson, Microzooplankton grazing of primary production at 140°W in the equatorial Pacific, *Deep Sea Res. Part II*, 43, 1227–1255, 1996.
- Yoder, J. A., S. G. Ackleson, R. T. Barber, P. Flament, and W. M. Balch, A line in the sea, *Nature*, 371, 689–692, 1994.
- Wei, K. Y., Z.-W. Zhang, M.-T. Chen, A. R. Isern, C.-H. Wang, and M. Leinen, Late Quaternary Paleoceanography of the central equatorial Pacific: A quantitative record of planktonic foraminiferal, isotopic, organic carbon and calcium carbonate changes, *J. Geol. Soc. China*, 37, 475–496, 1995.

E. Thomas and K. K. Turekian, Center for the Study of Global Change, Department of Geology and Geophysics, Yale University, P.O. Box 208109, New Haven, CT 06520-8109. (ethomas@wesleyan.edu)

K.-Y. Wei, Department of Geology, National Taiwan University, P.O. Box 23-13, Taipei 10617, Taiwan.

(Received October 27, 1998; revised January 28, 2000; accepted March 14, 2000.)